IASI Flux and Temperature

Plan de gestion de données créé à l'aide de DMP OPIDoR

Créateurs du PGD : Ana Parracho, Sarah Safieddine, Maya George

Affiliation du créateur principal : CNRS

Modèle du PGD : ERC DMP

Dernière modification du PGD: 25/01/2021

Financeur : Conseil européen de la recherche (European Research Council, ERC)

Numéro de subvention: 742909

Résumé du projet :

For the last few years, the record for the hottest month since modern meteorology observations began has been beaten several times. This, along with the recent so-called "hiatus" in the warming trend and the Paris climate agreement, has attracted scientific and public attention to the question of how reliable is the historical temperature record. Although the role of satellites in observing the variability and change of the Earth system has increased in recent decades, remotely-sensed observations are still underexploited to accurately assess climate change fingerprints. The IASI - Flux and Temperature (IASI-FT) project aims at providing new benchmarks for top-of-atmosphere radiative flux and temperature observations using the calibrated radiances measured twice a day at any location by the IASI instrument on the suite of MetOp satellites. The main challenge is to achieve the stringent accuracy and stability necessary for climate studies, particularly for climate trends. The development of innovative algorithms and statistical tools is proposed, to generate climate data records at the global scale, of (1) spectrally resolved outgoing radiances, (2) land and sea skin surface temperatures, and (3) temperatures at selected altitudes. Time series of these quantities are compared with in situ and other satellite observations, when available, atmospheric reanalyses, and climate model simulations. The observed trends are analyzed at seasonal and regional scales in order to disentangle natural (weather/dynamical) variability and human-induced climate forcings. This project, while clearly research-oriented, will lead towards an operational integrated observational strategy for the Earth climate system, given that the IASI program started in 2006 and will last until 2040 at least.

Chercheur Principal: Cathy Clerbaux

Identifiant ORCID: 0000-0003-0394-7200

Contact pour les Données : Maya George

Produits de recherche:

1. IASI SST : IASI-derived sea surface temperature (Jeu de données)

2. IASI SKIN TEMP: IASI SKIN TEMPERATURE (Jeu de données)

3. IASI OLR : IASI OUTGOING LONGWAVE RADIATION (Jeu de données)

IASI Flux and Temperature

Summary **Project Acronym** IASI-FT **Project Number** 742909 Provide a dataset summary (Several datasets may be included into a single DMP) Three products are developed within the IASI-FT project: IASI Sea Surface Temperature (SST), IASI Skin Temperature (SkT) and IASI Outgoing Longwave Radiation (OLR). 1) IASI SST is a monthly (L3), 1x1 degree global dataset of skin temperatures over the sea derived from the IASI satellite radiances data. The data was computed using Planck's law and simple atmospheric corrections. This method was applied to reprocessed IASI L1C data, in order to produce a homogeneous SST data record from 2008 to the present. The dataset consists of monthly mean TXT files of around 370 KB each (total 52,7 MB for the 2008-2019 period). Each file is a grid organized by latitude (-90 to 89 °N) and longitude (-180 to 180 °E). The land and sea ice grid points are filled using NaN. Unit: Kelvin. Separator: space. 2) IASI Skin Temperature is a monthly (L3), 1x1 degree global dataset of skin temperatures over land derived from the IASI satellite radiances data. The data was computed using artificial neural networks over a selected set of IASI radiance channels. trained with ERA5-skin temperature product. Emissivity is also used as input in the neural network (Safieddine et al., 2020). This method was applied to the whole IASI time series to produce a homogeneous skin temperature data record from Sept. 2007 to the The dataset consists of monthly mean TXT files of around 350 KB each (~total 52 MB for the 200709-201912 period). Each file is a grid organized by latitude (-90 to 89 °N) and longitude (-180 to 180 °E). Night and day data are separated based on the local time of IASI's overpass at each grid point. The sea grid points are filled using NaN. Unit: degree Celsius. Separator: comma.

3) The IASI spectrally resolved Outgoing Longwave Radiation product (IASI OLR) is a monthly (L3), 2x2 degree global dataset of spectral OLR derived from the clear-sky IASI satellite radiance measurements in the range 645-2300 cm-1 at the 0.25 cm-1 native spectral sampling of the L1C spectra. The algorithm for the conversion of the spectra to the OLR is detailed in Whitburn et al. (2020, Journal of Climate). It relies on precalculated angular distribution models (ADMs) constructed for over 140,000 different scene types (characterized by surface and atmospheric parameters). These were selected from one year of Copernicus

Atmosphere Monitoring Service (CAMS) reanalysis data. The method was applied to reprocessed IASI L1C data, in order to produce a homogeneous spectral OLR dataset from 2008 to present. The dataset consists of monthly mean NETCDF files of around 500 MB each (total 64.5 GB for the 2008-2017 period available to date). Each file includes the gridded spectral OLR calculated separately for day and nighttime IASI observations. Missing values are set at NaN. Size of the grids: 91 lines (from -90 to 90 °N), 181 columns (from -180 to 180 °E), 6621 aisles (between 645 and 2300 cm-1 by step of 0.25 cm-1). Unit: W • m-2 • cm-1.

FAIR data and resources

IASI SST: IASI-derived sea surface temperature

1. Making data findable

Data is available for download at: https://iasi-ft.eu/data-access/SST/ DOI: 10.21413/iasi-ft_metopa_sst_ 3_latmos-ulb 10.21413/iasi-ft_metopb_sst_ 3_latmos-ulb et 10.21413/iasi-ft_metopc_sst_ 3_latmos-ulb
2. Making data openly accessible
The gridded monthly mean SST dataset are available at https://iasi-ft.eu/data-access/SST/ Data is made available as TXT files and can therefore be accessed by anyone, and used in all data analysis software.
3. Making data interoperable
Question sans réponse.
4. Increase data reuse
The data will remain re-usable for the foreseeable future. No embargo is foreseen. Data is compared with other reference datsets.
5. Allocation of resources and data security
Question sans réponse.
IASI SKIN TEMP : IASI SKIN TEMPERATURE 1. Making data findable

Data is available for download at: https://iasi-ft.eu/data-access/SkT/

DOI: 10.21413/iasi-ft_metopa_skt_l3_latmos-ulb 10.21413/iasi-ft_metopb_skt_l3_latmos-ulb et_10.21413/iasi-ft_metopc_skt_l3_latmos-ulb

2. Making data openly accessible

The gridded monthly mean Skin Temperature dataset are available at https://iasi-ft.eu/data-access/SkT/Data is made available as TXT files and can therefore be accessed by anyone, and used in all data analysis software.
3. Making data interoperable
Question sans réponse.
4. Increase data reuse
The data will remain re-usable for the foreseeable future. No embargo is foreseen. Data is compared with other reference datsets.
5. Allocation of resources and data security
Question sans réponse.
IASI OLR : IASI OUTGOING LONGWAVE RADIATION
1. Making data findable
Data is available for download at: https://iasi-ft.eu/data-access/OLR/ DOI: 10.21413/iasi-ft_metopa_olr_l3_ulb-latmos
2. Making data openly accessible
The gridded monthly mean OLR dataset are available at https://iasi-ft.eu/data-access/OLR/Data is made available as NETCDF files.
3. Making data interoperable
Question sans réponse.

4. Increase data reuse	
The data will remain re-usable for the foreseeable future. No embargo is foreseen. Data is compared with other reference datsets.	
5. Allocation of resources and data security	
Question sans réponse.	